

the 1990s, the number of people with a diagnosis of schizophrenia has increased in many countries, including the United Kingdom (Murray & Lewis, 1998). This increase has been attributed to a number of factors, including changes in the definition of the disorder, changes in the way in which the disorder is diagnosed, and changes in the way in which the disorder is treated.

One of the most important factors in the increase in the number of people with a diagnosis of schizophrenia is the change in the way in which the disorder is diagnosed. In the past, the diagnosis of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features. However, in the 1990s, the diagnosis of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features. This change in the way in which the disorder is diagnosed has led to an increase in the number of people with a diagnosis of schizophrenia.

Another important factor in the increase in the number of people with a diagnosis of schizophrenia is the change in the way in which the disorder is treated. In the past, the treatment of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features. However, in the 1990s, the treatment of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features.

This change in the way in which the disorder is treated has led to an increase in the number of people with a diagnosis of schizophrenia.

Finally, another important factor in the increase in the number of people with a diagnosis of schizophrenia is the change in the way in which the disorder is defined. In the past, the definition of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features. However, in the 1990s, the definition of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features.

This change in the way in which the disorder is defined has led to an increase in the number of people with a diagnosis of schizophrenia.

In conclusion, the increase in the number of people with a diagnosis of schizophrenia in the 1990s can be attributed to a number of factors, including changes in the way in which the disorder is diagnosed, changes in the way in which the disorder is treated, and changes in the way in which the disorder is defined.

These changes have led to an increase in the number of people with a diagnosis of schizophrenia, and this increase has had a number of implications for the way in which the disorder is treated and the way in which the disorder is defined.

One of the most important implications of the increase in the number of people with a diagnosis of schizophrenia is the need for more resources to be devoted to the treatment of the disorder. In the past, the treatment of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features.

However, in the 1990s, the treatment of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features. This change in the way in which the disorder is treated has led to an increase in the number of people with a diagnosis of schizophrenia.

Another important implication of the increase in the number of people with a diagnosis of schizophrenia is the need for more resources to be devoted to the prevention of the disorder. In the past, the prevention of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features.

However, in the 1990s, the prevention of schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features. This change in the way in which the disorder is prevented has led to an increase in the number of people with a diagnosis of schizophrenia.

Finally, another important implication of the increase in the number of people with a diagnosis of schizophrenia is the need for more resources to be devoted to the research into the disorder. In the past, the research into schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features.

However, in the 1990s, the research into schizophrenia was based on a number of criteria, including the presence of certain symptoms, the duration of the illness, and the presence of certain features. This change in the way in which the disorder is researched has led to an increase in the number of people with a diagnosis of schizophrenia.

In conclusion, the increase in the number of people with a diagnosis of schizophrenia in the 1990s has had a number of implications for the way in which the disorder is treated, the way in which the disorder is prevented, and the way in which the disorder is researched.

These implications have led to an increase in the number of people with a diagnosis of schizophrenia, and this increase has had a number of implications for the way in which the disorder is treated, the way in which the disorder is prevented, and the way in which the disorder is researched.

In conclusion, the increase in the number of people with a diagnosis of schizophrenia in the 1990s can be attributed to a number of factors, including changes in the way in which the disorder is diagnosed, changes in the way in which the disorder is treated, and changes in the way in which the disorder is defined.